

AD-A170 649


MULTIPLICATIVE STOCHASTIC PROCESSES INVOLVING THE
TIME-DERIVATIVE OF A M.A. (U) STATE UNIV OF NEW YORK AT
BUFFALO DEPT OF CHEMISTRY M F ARNOLDUS ET AL. JUL 86
UBUFFALO/DC/86/TR-11 N00014-86-K-0043


1/1

UNCLASSIFIED

F/G 12/1

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

12

OFFICE OF NAVAL RESEARCH

Contract N00014-86-K-0043

TECHNICAL REPORT No. 11

Multiplicative Stochastic Processes Involving the Time-Derivative
of a Markov Process

by

Henk F. Arnoldus and Thomas F. George

Prepared for Publication

in

Journal of Mathematical Physics

Departments of Chemistry and Physics
State University of New York at Buffalo
Buffalo, New York 14260

July 1986

Reproduction in whole or in part is permitted for any purpose of the
United States Government.

This document has been approved for public release and sale;
its distribution is unlimited.



AD-A170 649

DTIC FILE COPY

86 8 11 009

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

AD-A170649

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE				
4. PERFORMING ORGANIZATION REPORT NUMBER(S) UBUFFALO/DC/86/TR-11			5. MONITORING ORGANIZATION REPORT NUMBER(S)	
6a. NAME OF PERFORMING ORGANIZATION Depts. Chemistry & Physics State University of New York		6b. OFFICE SYMBOL (If applicable)		7a. NAME OF MONITORING ORGANIZATION
6c. ADDRESS (City, State and ZIP Code) Fronczak Hall, Amherst Campus Buffalo, New York 14260			7b. ADDRESS (City, State and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable)		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Contract N00014-86-K-0043
8c. ADDRESS (City, State and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217			10. SOURCE OF FUNDING NOS.	
			PROGRAM ELEMENT NO.	PROJECT NO.
			TASK NO.	WORK UNIT NO.
11. TITLE Multiplicative Stochastic Processes Involving the Time-Derivative of a Markov Process				
12. PERSONAL AUTHOR(S) Henk F. Arnoldus and Thomas F. George				
13a. TYPE OF REPORT		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Yr., Mo., Day) July 1986
				15. PAGE COUNT 24
16. SUPPLEMENTARY NOTATION Prepared for publication in Journal of Mathematical Physics				
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB GR.	STOCHASTIC PROCESSES	
			MULTIPLICATIVE	
			MARKOV PROCESS	
			TIME-DERIVATIVE	
			ATOMIC RESPONSE	
			PHASE-LOCKED RADIATION	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) The characteristic functional of the derivative $\dot{\phi}(t)$ of a Markov process $\phi(t)$ and the related multiplicative process $\sigma(t)$, which obeys the stochastic differential equation $i\dot{\sigma}(t) = (A + \dot{\phi}(t)B)\sigma(t)$, have been studied. Exact equations for the marginal characteristic functional and the marginal average of $\sigma(t)$ are derived. The first equation is applied to obtain a set of equations for the marginal moments of $\dot{\phi}(t)$ in terms of the prescribed properties of $\phi(t)$. It is illustrated by an example how these equations can be solved, and it is shown in general that $\dot{\phi}(t)$ is delta-correlated, with a smooth background. The equation of motion for the marginal average of $\sigma(t)$ is solved for various cases, and it is shown how closed-form analytical expressions for the average $\langle \sigma(t) \rangle$ can be obtained.				
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT <input checked="" type="checkbox"/> DTIC USERS <input type="checkbox"/>			21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson			22b. TELEPHONE NUMBER (Include Area Code) (202) 696-4410	22c. OFFICE SYMBOL

MULTIPLICATIVE STOCHASTIC PROCESSES INVOLVING THE TIME-DERIVATIVE
OF A MARKOV PROCESS

Henk F. Arnoldus and Thomas F. George
Departments of Physics & Astronomy and Chemistry
239 Fronczak Hall
State University of New York at Buffalo
Buffalo, New York 14260

ABSTRACT

The characteristic functional of the derivative $\dot{\phi}(t)$ of a Markov process $\phi(t)$ and the related multiplicative process $\sigma(t)$, which obeys the stochastic differential equation $i\dot{\sigma}(t) = (A + \phi(t)B)\sigma(t)$, have been studied. Exact equations for the marginal characteristic functional and the marginal average of $\sigma(t)$ are derived. The first equation is applied to obtain a set of equations for the marginal moments of $\phi(t)$ in terms of the prescribed properties of $\phi(t)$. It is illustrated by an example how these equations can be solved, and it is shown in general that $\phi(t)$ is delta-correlated, with a smooth background. The equation of motion for the marginal average of $\sigma(t)$ is solved for various cases, and it is shown how closed-form analytical expressions for the average $\langle\sigma(t)\rangle$ can be obtained.

PACS: 2.50, 5.40



Accession For	
NTIS	CRA&I <input checked="" type="checkbox"/>
DTIC	TAB <input type="checkbox"/>
Unannounced <input type="checkbox"/>	
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

I. INTRODUCTION

The equation of motion for the density operator of an atom in a finite-bandwidth laser field or the equation for the regression of the atomic dipole correlations assumes the general form^{1,2}

$$i \frac{d\sigma}{dt} = (A + \dot{\phi}(t)B)\sigma, \quad (1.1)$$

where A and B are linear operators in Liouville space, which act on the Liouville vector $\sigma(t)$. Here $\phi(t)$ represents the laser phase, which is considered to be a real-valued stochastic process. The fluctuating phase broadens the laser line, but the atom responds to the instantaneous frequency shift $\dot{\phi}(t)$, which is the time-derivative of the laser phase.³ The process $\dot{\phi}(t)$ is again a stochastic process, and via Eq. (1.1) the state of the atom or the correlation functions $\sigma(t)$ become stochastic quantities. The issue in quantum optics is then to solve the multiplicative stochastic differential equation (1.1) for the average $\langle\sigma(t)\rangle$. The first solution was obtained by Fox,⁴ who assumed the process $\dot{\phi}(t)$ to be Gaussian white noise, which corresponds to a diffusive Gaussian phase $\phi(t)$ (the Wiener-Lévy process). This result was generalized to a Gaussian process $\dot{\phi}(t)$ with a finite correlation time and an exponentially decaying correlation function⁵⁻⁷ (the Ornstein-Uhlenbeck process), and to a process $\dot{\phi}(t)$, which is again diffusive, but not Gaussian^{8,9} (the independent-increment process). Furthermore, Eq. (1.1) can be solved for $\langle\sigma(t)\rangle$ if we have $\dot{\phi}(t)$ as a Markov random-jump process,¹⁰⁻¹³ which models a multimode laser.^{14,15}

In these examples the solvability of the problem relies on the Gaussian property of $\dot{\phi}(t)$, or hinges on the prescribed stochastics of $\dot{\phi}(t)$. This implies that the process $\dot{\phi}(t)$ is actually considered to be the driving

process. For a single-mode laser in general, however, the phase fluctuations $\phi(t)$ are specified rather than the derivative $\dot{\phi}(t)$ of this process. A prime example would be the atomic response to phase-locked radiation,¹⁶ as it is generated for instance by some ring lasers.¹⁷ In this paper we shall develop a general method to solve Eq. (1.1) for the case that $\phi(t)$ is a given Markov process. The formal theory will be exemplified by a specific choice for $\phi(t)$, which models phase-locked radiation. Furthermore, we shall study the time-derivative of $\phi(t)$ itself and extract the stochastics of $\dot{\phi}(t)$ from the properties of $\phi(t)$.

II. THE STOCHASTICS OF $\phi(t)$

Let us define the phase $\phi(t)$ as a homogeneous Markov process.¹⁸ Then its stochastics is fixed by the probability distribution $P(\phi, t)$ and the conditional probability distribution $P_\tau(\phi_2 | \phi_1)$ ($\tau \geq 0$), which has the significance of the probability density for the occurrence of $\phi(t+\tau) = \phi_2$ if $\phi(t) = \phi_1$. For a homogeneous process this is independent of t by definition. The higher-order statistics is now determined by the Markov property.¹⁹ From the obvious relation

$$\int d\phi' P_{t-t_0}(\phi | \phi') P(\phi', t_0) = P(\phi, t), \quad t \geq t_0 \quad (2.1)$$

it follows that it is sufficient to prescribe the probability distribution $P(\phi, t)$ for a single time-point t_0 only. The time-evolution towards $t > t_0$ can then be found from Eq. (2.1) and $P_{t-t_0}(\phi | \phi')$.

The conditional probability distribution obeys the Master equation¹⁸

$$\frac{\partial}{\partial \tau} P_\tau(\phi_3 | \phi_1) = \int d\phi_2 \{W(\phi_3 | \phi_2) - a(\phi_2)\delta(\phi_3 - \phi_2)\} P_\tau(\phi_2 | \phi_1) \quad (2.2)$$

with $W(\phi' | \phi) \geq 0$ as the transition rate of the process from ϕ to ϕ' and

$$a(\phi) = \int d\phi' W(\phi'|\phi) , \quad (2.3)$$

which is the loss-rate of ϕ , independent of the final value ϕ' . The initial condition for Eq. (2.2) reads

$$P_0(\phi_3|\phi_1) = \delta(\phi_3 - \phi_1) , \quad (2.4)$$

so a given $W(\phi'|\phi)$ determines $P_\tau(\phi_3|\phi_1)$ for every $\tau \geq 0$. Hence the stochastics of a homogeneous Markov process $\phi(t)$ is fixed as soon as $P(\phi, t_0)$ and $W(\phi'|\phi)$ are prescribed. These functions will from now on be assumed to be given.

III. THE CHARACTERISTIC FUNCTIONAL

A convenient way to represent the stochastic properties of a stochastic process is by means of its characteristic functional.^{9,20} Since we are concerned with the process $\dot{\phi}(t)$, we define

$$Z_t[k] = \langle \exp(-i \int_{t_0}^t ds \dot{\phi}(s) k(s)) \rangle , \quad t \geq t_0 , \quad (3.1)$$

which is a functional of the test function $k(t)$. Here the angle brackets denote an average over the stochastic process $\phi(t)$ or $\dot{\phi}(t)$, whatever is prescribed. A general method to evaluate $Z_t[k]$ for the case where $\dot{\phi}(t)$ is a homogeneous Markov process has been given by van Kampen.²¹

Knowledge of the characteristic functional $Z_t[k]$ determines completely the stochastics of $\dot{\phi}(t)$, which can be seen as follows. Choose $k(s)$ as the sequence of δ -functions

$$k(s) = - \sum_{\ell=1}^n \delta(s - t_\ell) k_\ell , \quad t_\ell > t_0 , \quad (3.2)$$

and take $t = \infty$ in (3.1). Then we find

$$Z_{\infty}[k] = \langle \exp(ik_n \dot{\phi}(t_n) + \dots + ik_1 \dot{\phi}(t_1)) \rangle, \quad (3.3)$$

which is the moment-generating function of $\dot{\phi}(t)$. If we write $z_n(k_n, t_n; \dots; k_1, t_1)$ for $Z_{\infty}[k]$, then we can obtain the moments of $\psi(t) \equiv \dot{\phi}(t)$ according to

$$\langle \psi(t_n) \dots \psi(t_1) \rangle = (-i)^n \frac{\partial}{\partial k_n} \dots \frac{\partial}{\partial k_1} z_n(k_n, t_n; \dots; k_1, t_1) \Big|_{k_n = \dots = k_1 = 0}, \quad (3.4)$$

and the probability distributions by

$$\begin{aligned} \bar{P}_n(\psi_n, t_n; \dots; \psi_1, t_1) &= \frac{1}{(2\pi)^n} \int dk_n \dots dk_1 \\ &\times e^{-ik_n \psi_n - \dots - ik_1 \psi_1} z_n(k_n, t_n; \dots; k_1, t_1), \end{aligned} \quad (3.5)$$

where we have introduced \bar{P}_n in order to distinguish from the probability distributions for $\phi(t)$ itself.

IV. THE MARGINAL AVERAGE

A. GENERAL

The exponential in Eq. (3.1) is a functional of both $k(t)$ and $\dot{\phi}(t)$, so it depends on the values of $\dot{\phi}(t)$ in the complete interval $[t_0, t]$. After the average has been taken it will be only a functional of $k(t)$. The general attempt to evaluate averages of a functional is to derive an equation for the average. For subsequently solving this equation for functionals which involve Markov processes, this scheme is most conveniently carried out by an intermediate introduction of Burshtein's marginal averages.²² Since in our problem the stochastics of $\phi(t)$ is assumed to be given, the appropriate marginal characteristic functional, which is related to $Z_t[k]$, should be defined as

$$Q_t[\phi_0, k] = \langle \delta(\phi(t) - \phi_0) \exp(-i \int_{t_0}^t \dot{\phi}(s) k(s)) \rangle, \quad (4.1)$$

for $t \geq t_0$. The initial value is then

$$Q_{t_0}[\phi_0, k] = \langle \delta(\phi(t_0) - \phi_0) \rangle = P(\phi_0, t_0), \quad (4.2)$$

and $Z_t[k]$ follows from $Q_t[\phi_0, k]$ according to

$$Z_t[k] = \int d\phi_0 Q_t[\phi_0, k]. \quad (4.3)$$

For $t = t_0$ we find with Eq. (4.2) $Z_{t_0}[k] = \int d\phi_0 P(\phi_0, t_0) = 1$, in agreement with Eq. (3.1).

In order to derive an equation for the time-evolution of $Q_t[\phi_0, k]$, we first increase t by a small amount $\Delta t > 0$. This gives

$$\begin{aligned} Q_{t+\Delta t}[\phi_0, k] &= \langle \delta(\phi(t+\Delta t) - \phi_0) \exp\{-i(\phi(t+\Delta t) - \phi(t))k(t)\} \\ &\quad \times \exp(-i \int_{t_0}^t \dot{\phi}(s) k(s)) \rangle. \end{aligned} \quad (4.4)$$

Subsequently, we expand the exponential functional of $\dot{\phi}(s)$ in a series, and we take the average in (4.4) term by term. Hereafter, we apply the Master equation (2.2) for $P_{t+\Delta t}(\phi|\phi_0)$ and take the limit $\Delta t \rightarrow 0$. This yields an equation for the marginal average, and explicitly we find

$$\begin{aligned} \frac{\partial}{\partial t} Q_t[\phi_0, k] &= \int d\phi \{W(\phi_0|\phi) - a(\phi)\delta(\phi_0 - \phi)\} \\ &\quad \times e^{-i(\phi_0 - \phi)k(t)} Q_t[\phi, k]. \end{aligned} \quad (4.5)$$

The Markov process $\phi(t)$ is characterized by $P(\phi_0, t_0)$ and $W(\phi_0|\phi)$, which respectively determine the initial value and the time-evolution of $Q_t[\phi_0, k]$.

For a specific choice of $W(\phi_0|\phi)$, we have to solve Eq. (4.5), after which the characteristic functional $Z_t[k]$ can be obtained from Eq. (4.3).

Notice the resemblance between the result (4.5) and the Master equation (2.2). If we multiply Eq. (2.2) by $P(\phi_1, t_0)$, take $\tau = t - t_0$ and apply the relation (2.1), we find

$$\frac{\partial}{\partial t} P(\phi_0, t) = \int d\phi \{W(\phi_0|\phi) - a(\phi)\delta(\phi_0 - \phi)\} P(\phi, t) , \quad (4.6)$$

which is the Master equation for $P(\phi_0, t)$. This equation is identical to Eq. (4.5), including the initial condition (4.2), if we set $k(t) \equiv 0$. On the other hand, it follows from Eq. (4.1) that $Q_t[\phi_0, k] = \langle \delta(\phi(t) - \phi_0) \rangle = P(\phi_0, t)$ if we take $k(t) = 0$, so that in this case Eq. (4.5) should indeed reduce to Eq. (4.6).

B. INDEPENDENT INCREMENTS

In order to display the usefulness and applicability of the marginal-functional approach, we consider an example. Let us specify the transition rate by

$$W(\phi_0|\phi) = \gamma w(\phi_0 - \phi) , \quad \gamma > 0 , \quad (4.7)$$

where the function $w(\eta)$ is normalized as

$$\int d\eta w(\eta) = 1 . \quad (4.8)$$

The stochastic process $\phi(t)$ will be defined on the real axis, with $-\infty < \phi < \infty$. The assertion (4.7) states that the probability for a transition $\phi \rightarrow \phi_0$ depends only on the phase difference $\phi_0 - \phi$, and from Eq. (2.3) we find that $a(\phi) = \gamma$, so that the total loss-rate for ϕ is independent of ϕ . This is a diffusion process, and it is commonly referred to as the independent-increment process. As an initial condition for the probability

distribution, we take

$$P(\phi, t_0) = \delta(\phi) . \quad (4.9)$$

Comparison of the Master equations for $P_\tau(\phi|\phi')$ and $P(\phi, t)$ then shows that the probability distribution and the conditional probability distribution are related according to

$$P_{t-t_0}(\phi|\phi_0) = P(\phi-\phi_0, t) . \quad (4.10)$$

The Master equation (4.6) for $P(\phi, t)$ can be solved by Fourier transformation with respect to ϕ . If we write

$$\hat{P}(\rho, t) = \langle e^{i\rho\phi(t)} \rangle = \int_{-\infty}^{\infty} d\phi e^{i\rho\phi} P(\phi, t) , \quad (4.11)$$

which has $\hat{P}(\rho, t_0) = 1$ as the initial condition, then the solution of Eq. (4.6) is immediately seen to be

$$\hat{P}(\rho, t) = e^{\gamma(\hat{w}(\rho)-1)(t-t_0)} , \quad t \geq t_0 , \quad (4.12)$$

in terms of the Fourier transform $\hat{w}(\rho)$ of $w(\phi)$. Note that $\hat{w}(0) = 1$, as a result of the normalization (4.7). Along the very same lines we can solve Eq. (4.5) for the Fourier transform $\hat{Q}_t[\rho, k]$. We obtain

$$\hat{Q}_t[\rho, k] = \exp\left(-\gamma \int_{t_0}^t ds \int_{-\infty}^{\infty} d\phi (1 - e^{i\phi(\rho - k(s))}) w(\phi)\right) , \quad (4.13)$$

after which the characteristic functional follows from

$$Z_t[k] = \hat{Q}_t[0, k] , \quad (4.14)$$

which yields the familiar result .⁸

V. THE MARGINAL MOMENTS

A. GENERAL

If we take $k(s)$ as the sequence of delta functions (3.2) in the definition (4.1) of the marginal characteristic functional, it assumes the form

$$Q_t[\phi_0, k] = \langle \delta(\phi(t) - \phi_0) \exp(i \sum_{\ell=1}^n k_\ell \psi(t_\ell) \theta(t - t_\ell)) \rangle, \quad (5.1)$$

with $\psi(t) = \dot{\phi}(t)$ and $\theta(t)$ the unit-step function. Just as we can find the moments $\langle \psi(t_n) \dots \psi(t_1) \rangle$ of $\psi(t)$ from $Z_\infty[k]$, we can obtain the marginal moments $\langle \delta(\phi(t) - \phi_0) \psi(t_n) \dots \psi(t_1) \rangle$ from $Q_t[\phi_0, k]$. Obviously the integral over ϕ_0 of the marginal moments yields the moments. The characteristic functional $Z_t[k]$ becomes independent of t if $t > t_\ell$ for all ℓ , but $Q_t[\phi_0, k]$ remains time-dependent. This is due to the appearance of $\delta(\phi(t) - \phi_0)$. Furthermore, the time t is a dynamical variable in Eq. (4.5), so that care should be exercised in the time-ordering. The marginal moments follow from $Q_t[\phi_0, k]$ by differentiation, according to

$$\begin{aligned} \langle \delta(\phi(t) - \phi_0) \psi(t_n) \dots \psi(t_1) \rangle \theta(t - t_n) \dots \theta(t - t_1) = \\ (-i)^n \frac{\partial}{\partial k_n} \dots \frac{\partial}{\partial k_1} Q_t[\phi_0, k] \Big|_{k_n = \dots = k_1 = 0}. \end{aligned} \quad (5.2)$$

Equation (4.5) for $Q_t[\phi_0, k]$ implies an equation for the marginal moments. First, we note that

$$\begin{aligned} \exp\{-i(\phi_0 - \phi)k(t)\} Q_t[\phi, k] = \\ \langle \delta(\phi(t) - \phi) \exp(i \sum_{\ell=1}^n k_\ell \{(\phi_0 - \phi)\delta(t - t_\ell) + \psi(t_\ell)\theta(t - t_\ell)\}) \rangle. \end{aligned} \quad (5.3)$$

After substituting this expression in the right-hand side of Eq. (4.5), differentiating with respect to k_n, \dots, k_1 , setting $k_n = \dots = k_1 = 0$ and integrating over time, we obtain

$$\begin{aligned} \langle \delta(\phi(t) - \phi_0) \psi(t_n) \dots \psi(t_1) \rangle \theta(t - t_n) \dots \theta(t - t_1) &= \int d\phi \{W(\phi_0 | \phi) - a(\phi) \delta(\phi_0 - \phi)\} \\ &\times \int_{t_0}^t dt' \langle \delta(\phi(t') - \phi) \{(\phi_0 - \phi) \delta(t' - t_n) + \psi(t_n) \theta(t' - t_n)\} \\ &\dots \{(\phi_0 - \phi) \delta(t' - t_1) + \psi(t_1) \theta(t' - t_1)\} \rangle. \end{aligned} \quad (5.4)$$

When we set $t > t_\ell$ for all ℓ , we have a Master-like equation for $\langle \delta(\phi(t) - \phi_0) \psi(t_n) \dots \psi(t_1) \rangle$, and the lower-order marginal moments $\langle \delta(\phi(t) - \phi_0) \psi(t_m) \dots \psi(t_1) \rangle$ with $m < n$ appear as inhomogeneous terms. Hence, Eq. (5.4) should be solved successively for $n = 1, n = 2, \dots$. We note that Eq. (5.4) provides an explicit expression for $\langle \psi(t_n) \dots \psi(t_1) \rangle$ in terms of the lower-order marginal moments after an integration over ϕ_0 . Indeed, from the property

$$\int d\phi_0 \{W(\phi_0 | \phi) - a(\phi) \delta(\phi_0 - \phi)\} = 0, \quad (5.5)$$

the term with $\langle \delta(\phi(t') - \phi) \psi(t_n) \dots \psi(t_1) \rangle$ on the right-hand side of Eq. (5.4) vanishes after an integration over ϕ_0 .

B. LOWEST ORDERS

In order to exhibit clearly the structure of the equation for the marginal moments, we consider the cases $n = 1$ and $n = 2$ in some more detail. After a slight rearrangement, Eq. (5.4) for $n = 1$ can be written as

$$\begin{aligned} \langle \delta(\phi(t) - \phi_0) \psi(t_1) \rangle &= \int d\phi \{W(\phi_0 | \phi) - a(\phi) \delta(\phi_0 - \phi)\} \\ &\times \{(\phi_0 - \phi) P(\phi, t_1) + \int_{t_1}^t dt' \langle \delta(\phi(t') - \phi) \psi(t_1) \rangle\}, \end{aligned} \quad (5.6)$$

for $t \geq t_1$. This integral equation in time is equivalent to the differential equation

$$\begin{aligned} \frac{\partial}{\partial t} \langle \delta(\phi(t) - \phi_0) \psi(t_1) \rangle &= \int d\phi \{ W(\phi_0 | \phi) - a(\phi) \delta(\phi_0 - \phi) \} \\ &\times \langle \delta(\phi(t) - \phi) \psi(t_1) \rangle, \end{aligned} \quad (5.7)$$

together with the initial condition

$$\langle \delta(\phi(t_1) - \phi_0) \psi(t_1) \rangle = \int d\phi \{ W(\phi_0 | \phi) - a(\phi) \delta(\phi_0 - \phi) \} (\phi_0 - \phi) P(\phi, t_1). \quad (5.8)$$

The equation for the first marginal average $\langle \delta(\phi(t) - \phi_0) \psi(t_1) \rangle$ is identical to the Master equation (2.2), but with a different initial value.

Integration of (5.8) over ϕ_0 yields

$$\langle \psi(t_1) \rangle = \int d\phi \int d\phi_0 W(\phi_0 | \phi) (\phi_0 - \phi) P(\phi, t_1), \quad (5.9)$$

which expresses explicitly the average of $\langle \psi(t_1) \rangle$ in the given functions $W(\phi_0 | \phi)$ and $P(\phi, t_1)$. With the aid of the Master equation, we can cast (5.9) in the form

$$\begin{aligned} \langle \psi(t_1) \rangle &= \int d\phi \phi \frac{\partial}{\partial t_1} P(\phi, t_1) \\ &= \frac{d}{dt_1} \langle \phi(t_1) \rangle, \end{aligned} \quad (5.10)$$

as it should be.

The solution of Eq. (5.6) for $\langle \delta(\phi(t) - \phi_0) \psi(t_1) \rangle$ provides the input for the explicit expression for the two-time correlation function, which becomes

$$\begin{aligned} \langle \psi(t_2) \psi(t_1) \rangle &= \int d\phi \int d\phi_0 \{ W(\phi_0 | \phi) - a(\phi) \delta(\phi_0 - \phi) \} \\ &\times \{ (\phi_0 - \phi)^2 \delta(t_2 - t_1) P(\phi, t_2) \\ &+ (\phi_0 - \phi) (\langle \delta(\phi(t_2) - \phi) \psi(t_1) \rangle \theta(t_2 - t_1) \\ &+ \langle \delta(\phi(t_1) - \phi) \psi(t_2) \rangle \theta(t_1 - t_2)) \}. \end{aligned} \quad (5.11)$$

The appearance of $\delta(t_2 - t_1)$ shows that the time-derivative of any Markov process is δ -correlated with a continuous background.

C. RANDOM JUMPS

Equation (5.11) for instance might seem awkward, but it is really straightforward in its application. Let us illustrate this with an example. Consider the random-jump process $\phi(t)$, defined as a stationary process with transition rate

$$W(\phi|\phi') = \gamma P(\phi) \quad , \quad \gamma > 0 \quad , \quad (5.12)$$

in terms of an arbitrary probability distribution $P(\phi)$. Eq. (5.12) is equivalent to the statement that the probability for a transition $\phi' \rightarrow \phi$ is independent of the initial value ϕ' .¹³ From Eq. (5.9) we immediately derive

$$\langle \psi(t_1) \rangle = 0 \quad , \quad (5.13)$$

which is, in view of (5.10), necessary for a stationary process. From (2.3) we obtain $a(\phi) = \gamma$, and the solution of Eq. (5.7), with initial value (5.8), is readily found to be

$$\langle \delta(\phi(t) - \phi_0) \psi(t_1) \rangle = \gamma P(\phi_0) (\phi_0 - b_1) e^{-\gamma(t-t_1)} \quad , \quad t \geq t_1 \quad . \quad (5.14)$$

Here we have introduced the moments of $P(\phi)$ as

$$b_n = \int d\phi \phi^n P(\phi) \quad , \quad (5.15)$$

which are parameters of the process $\phi(t)$. Solution (5.14) can be substituted into Eq. (5.11), which gives the correlation function

$$\langle \psi(t_1) \psi(t_2) \rangle = \gamma (b_2 - b_1^2) \{ 2\delta(t_1 - t_2) - \gamma e^{-\gamma|t_2 - t_1|} \} \quad , \quad (5.16)$$

for all t_1, t_2 . From (5.15) it follows that

$$b_2 - b_1^2 \geq 0, \quad (5.17)$$

so that for $t_1 \neq t_2$ the correlation (5.16) is negative. For $t_1 = t_2$ the δ -function dominates the negative term, so that $\langle \psi(t_1)^2 \rangle$ is positive, as it should be.

VI. THE MULTIPLICATIVE PROCESS

So far we have considered the stochastics of $\dot{\phi}(t)$ itself, and its characteristic functional. In this section we shall generalize the method, in order to solve the multiplicative equation (1.1). To this end we write the formal solution of (1.1) for the stochastic vector $\sigma(t)$ as

$$\sigma(t) = e^{-iA(t-t_0)} T \exp\left[-i \int_{t_0}^t ds \dot{\phi}(s) \tilde{B}(s)\right] \sigma(t_0), \quad (6.1)$$

where T is the time-ordering operator and $\tilde{B}(t)$ is defined as

$$\tilde{B}(t) = e^{iA(t-t_0)} B e^{-iA(t-t_0)}. \quad (6.2)$$

In close analogy to the definition of $Q_t[\phi_0, k]$ in Eq. (4.1), we now introduce the marginal average of $\sigma(t)$ by

$$\zeta(\phi_0, t) = \langle \delta(\phi(t) - \phi_0) \sigma(t) \rangle. \quad (6.3)$$

Then we substitute the expression (6.1) for $\sigma(t)$ and replace t by $t + \Delta t$, which gives a similar formula as Eq. (4.4). That this can also be done for the time-ordered exponential is sometimes referred to as the semi-group property of the evolution operator. Along the same lines that led to Eq. (4.5) we now find

$$\begin{aligned} i \frac{\partial}{\partial t} \zeta(\phi_0, t) = & A \zeta(\phi_0, t) \\ & + i \int d\phi \{ W(\phi_0 | \phi) - a(\phi) \delta(\phi_0 - \phi) \} e^{-i(\phi_0 - \phi)B} \zeta(\phi, t), \end{aligned} \quad (6.4)$$

or equivalently

$$\left(i\frac{\partial}{\partial t} - A + ia(\phi_0)\right) \zeta(\phi_0, t) = i \int d\phi W(\phi_0|\phi) e^{-i(\phi_0 - \phi)B} \zeta(\phi, t) . \quad (6.5)$$

Notice that the operator B appears in the exponential, rather than $\tilde{B}(t)$, as could be expected by analogy with the characteristic functional. For a given stochastic process $\phi(t)$, e.g., a given $W(\phi|\phi')$ and $P(\phi, t_0)$, we have to solve Eq. (6.4) with the initial condition

$$\zeta(\phi_0, t_0) = \langle \delta(\phi(t_0) - \phi_0) \sigma(t_0) \rangle , \quad (6.6)$$

after which $\langle \sigma(t) \rangle$ follows from

$$\langle \sigma(t) \rangle = \int d\phi_0 \zeta(\phi_0, t) . \quad (6.7)$$

For a given non-stochastic state $\sigma(t_0)$, the initial condition reduces to

$$\zeta(\phi_0, t_0) = P(\phi_0, t_0) \sigma(t_0) , \quad (6.8)$$

which differs from (6.6) by the fact that there are no initial correlations. This means that the process $\sigma(t)$ has no memory to times smaller than t_0 , and consequently its evolution for $t \geq t_0$ is completely determined by its initial state $\sigma(t_0)$. It was emphasized by Arnoldus and Nienhuis¹³ that the common choice $\zeta(\phi_0, t_0) = P(\phi_0, t_0) \langle \sigma(t_0) \rangle$ is merely an approximation which only holds for small correlation times of $\phi(t)$.

VII. SOLUTIONS

A. INDEPENDENT INCREMENTS

Equation (6.5) for the marginal average of $\sigma(t)$ can be solved for the independent-increment process with the same procedure as in Section IV, where we obtained the characteristic functional. If we adopt the Fourier transform

$$\tilde{\zeta}(\rho, t) = \int_{-\infty}^{\infty} d\phi e^{i\rho\phi} \zeta(\phi, t) = \langle e^{i\rho\phi(t)} \sigma(t) \rangle, \quad (7.1)$$

where the second equality follows after application of Eq. (6.3), then

$\langle \sigma(t) \rangle$ can be found from

$$\langle \sigma(t) \rangle = \tilde{\zeta}(0, t). \quad (7.2)$$

With the technique of Section IV we can find $\tilde{\zeta}(\rho, t)$, and if we differentiate the result with respect to time, we find

$$i \frac{\partial}{\partial t} \tilde{\zeta}(\rho, t) = (A - i\hat{W}(\rho)) \tilde{\zeta}(\rho, t), \quad (7.3)$$

with

$$\hat{W}(\rho) = \gamma \int_{-\infty}^{\infty} d\eta (1 - e^{i\eta(\rho-B)}) w(\eta). \quad (7.4)$$

The operator $\hat{W}(\rho)$ accounts for the phase fluctuations. If we set $\rho = 0$ in Eq. (7.3), we achieve the equation for $\langle \sigma(t) \rangle$, with solution

$$\langle \sigma(t) \rangle = e^{-i(A - i\hat{W}(0))(t-t_0)} \langle \sigma(t_0) \rangle, \quad (7.5)$$

for $t \geq t_0$. We note that $\langle \sigma(t) \rangle$ can be expressed in terms of $\langle \sigma(t_0) \rangle$ for this process, so that there are no initial correlations for the diffusion process. The process $\dot{\phi}(t)$ has no memory, and with the results of Section V it can be shown that $\dot{\phi}(t)$ is indeed delta-correlated. This means that $\langle \dot{\phi}(t_n) \dots \dot{\phi}(t_1) \rangle$ for all n contains only δ -functions, which implies the factorization in (7.5).

A special case arises if we take

$$\gamma w(\eta) = \gamma \delta(\eta) + \lambda \delta''(\eta), \quad \lambda > 0, \quad (7.6)$$

where the primes on the δ -function denote differentiation with respect to its argument. It is easy to check that this process is the Wiener-Lévy process, or the phase-diffusion process. If we substitute (7.6) into (4.12), we find that $P(\phi, t)$ is Gaussian, and obviously this is the only Gaussian limit of the diffusion process. The operator $\hat{W}(\rho)$ in (7.4) reduces to

$$\hat{W}(\rho) = \lambda(\rho - B)^2, \quad (7.7)$$

and the equation for $\langle \sigma(t) \rangle$ becomes

$$i \frac{d}{dt} \langle \sigma(t) \rangle = (A - i\lambda B^2) \langle \sigma(t) \rangle, \quad (7.8)$$

which is the result of Fox⁴.

B. ORNSTEIN-UHLENBECK PROCESS

The diffusion process has no memory and is essentially non-stationary. The initial distribution $P(\phi, t_0) = \delta(\phi)$ diffuses over the whole ϕ -axis, $-\infty < \phi < \infty$. The inclusion of a finite memory-time can stabilize this process. Let us define the transition probability as

$$W(\phi_0 | \phi) - a(\phi)\delta(\phi_0 - \phi) = \lambda \delta''(\phi_0 - \phi) + \gamma \phi \delta'(\phi_0 - \phi), \quad \begin{matrix} \lambda > 0 \\ \gamma > 0 \end{matrix}, \quad (7.9)$$

Then the Master equation (4.6) for $P(\phi, t)$ becomes the Fokker-Planck equation¹⁸

$$\frac{\partial}{\partial t} P(\phi, t) = \left(\lambda \frac{\partial^2}{\partial \phi^2} + \gamma \frac{\partial}{\partial \phi} \phi \right) P(\phi, t), \quad (7.10)$$

which has the solution for $t \rightarrow \infty$

$$P(\phi) = (2\pi\sigma^2)^{-\frac{1}{2}} e^{-\phi^2/2\sigma^2}, \quad \sigma^2 = \lambda/\gamma. \quad (7.11)$$

This $P(\phi)$, together with $W(\phi_0|\phi)$ from (7.9), defines a stationary Gaussian Markov process, the Ornstein-Uhlenbeck process. In the limit $\gamma \rightarrow 0$ and λ finite (so $\sigma^2 \rightarrow \infty$), the process $\phi(t)$ reduces to the Wiener-Lévy process from the previous paragraph. From (7.11) we see that $\phi(t)$ is centered around $\phi = 0$. The distribution is Gaussian with a variance σ^2 around the average $\phi = 0$. The preference for $\phi = 0$ expresses that this process can be considered as a model for phase-locked radiation.

With the specific choice (7.9) for the transition rate, the Master equation (6.5) assumes the form of a second-order partial differential equation. We obtain

$$\left(i\frac{\partial}{\partial t} - A + i\lambda B^2\right) \zeta(\phi, t) = i\gamma\left\{\sigma^2\left(2iB + \frac{\partial}{\partial \phi}\right) \frac{\partial}{\partial \phi} + \left(iB + \frac{\partial}{\partial \phi}\right)\phi\right\} \zeta(\phi, t) . \quad (7.12)$$

In the limit $\gamma \rightarrow 0$ and $\lambda = \gamma\sigma^2$ finite, we recover (the Fourier inverse of) Eq. (7.3) with $\hat{W}(\rho)$ from Eq. (7.7).

In order to obtain a solution of Eq. (7.12), we start with a Fourier transform with respect to ϕ . The transformed equation then reads

$$\left(i\frac{\partial}{\partial t} - A + i\lambda B^2\right) \tilde{\zeta}(\rho, t) = -i\gamma\left\{\sigma^2(\rho - 2B)\rho + (\rho - B) \frac{\partial}{\partial \rho}\right\} \tilde{\zeta}(\rho, t) , \quad (7.13)$$

which is still a partial differential equation. Since we are interested in $\tilde{\zeta}(0, t) = \langle \sigma(t) \rangle$, the obvious approach²³ would be a Taylor expansion around $\rho = 0$. This yields however a cumbersome inhomogeneous four-term recurrence relation for the Taylor coefficients. This can be avoided by the transformation¹⁵

$$\tilde{\zeta}(\rho, t) = \hat{P}(\rho) \hat{g}(\rho, t) , \quad (7.14)$$

which defines $\hat{g}(\rho, t)$. The Fourier transform of the probability distribution is explicitly

$$\hat{P}(\rho) = \langle e^{i\rho\phi(t)} \rangle = e^{-\frac{1}{2}\sigma^2\rho^2}, \quad (7.15)$$

and in particular we have $\hat{P}(0) = 1$. The equation for $\hat{g}(\rho, t)$ becomes

$$(i\frac{\partial}{\partial t} - A + i\lambda B^2)\hat{g}(\rho, t) = -i\gamma\{\rho\frac{\partial}{\partial\rho} - B(\sigma^2\rho + \frac{\partial}{\partial\rho})\}\hat{g}(\rho, t), \quad (7.16)$$

and it has to be solved for

$$\hat{g}(0, t) = \langle \sigma(t) \rangle. \quad (7.17)$$

Let us define the Taylor coefficients $\pi_n(t)$ by the expansion

$$\hat{g}(\rho, t) = \sum_{n=0}^{\infty} \frac{(i\rho)^n}{n!} \pi_n(t), \quad (7.18)$$

which can be inverted as

$$\pi_n(t) = \langle \sigma(t) \left(\frac{\partial}{\partial i\rho} \right)^n e^{i\rho\phi(t) - \frac{1}{2}(i\rho\sigma)^2} \rangle_{\rho=0}. \quad (7.19)$$

Substitution of (7.18) into (7.16) then gives the equation for the Taylor coefficients

$$(i\frac{\partial}{\partial t} - A + i\lambda B^2 + i\gamma n)\pi_n(t) = \gamma B(n\sigma^2 \pi_{n-1}(t) - \pi_{n+1}(t)), \quad (7.20).$$

which has to be solved for

$$\pi_0(t) = \langle \sigma(t) \rangle. \quad (7.21)$$

Equation (7.20) looks like a homogeneous three-term recurrence relation, but it will be shown below that the time-derivative $\partial/\partial t$ gives rise to an inhomogeneous contribution. Notice that for $n = 0$ Eq.(7.20) reduces to a two-term relation between $\pi_0(t)$ and $\pi_1(t)$ only.

Eq. (7.20) is most easily solved in the Laplace domain. If we introduce

$$\tilde{\pi}_n(\omega) = \int_{t_0}^{\infty} dt e^{i\omega(t-t_0)} \pi_n(t) , \quad (7.22)$$

then (7.20) attains the form

$$(\omega - A + i\lambda B^2 + i\gamma n)\tilde{\pi}(\omega) - \gamma B(n\sigma^2 \tilde{\pi}_{n-1}(\omega) - \tilde{\pi}_{n+1}(\omega)) = i\pi_n(t_0) . \quad (7.23)$$

Here the initial values $\pi_n(t_0)$ for $n = 0, 1, 2, \dots$ appear as inhomogeneous terms. The set $\pi_n(t_0)$ for all n represents the initial correlations of $\sigma(t)$ on $t = t_0$, and they connect the time-evolution of $\langle \sigma(t) \rangle$ for $t > t_0$ to its recent past.¹³ In other words, Eq. (7.23) relates the set $\pi_n(t)$ for $t > t_0$ to the initial set $\pi_n(t_0)$.

Equation (7.23) can be solved for an arbitrary initial set $\pi_n(t_0)$ by standard techniques,²⁴ but the solution is not transparent. In order to elucidate the structure of the solution, we assume a non-stochastic initial state $\sigma(t_0)$. From Eq. (7.1) we then find at $t = t_0$

$$\hat{\zeta}(\rho, t_0) = \hat{P}(\rho) \sigma(t_0) , \quad (7.24)$$

and from Eq. (7.14) we obtain

$$\hat{g}(\rho, t_0) = \sigma(t_0) . \quad (7.25)$$

Hence at $t = t_0$ the vector $\hat{g}(\rho, t_0)$ is independent of ρ , and therefore the expansion coefficients are simply

$$\pi_n(t_0) = \delta_{n,0} \sigma(t_0) . \quad (7.26)$$

Then only the recurrence relation for $n = 0$ is inhomogeneous, and the solution of (7.23) for $\tilde{\pi}_0(\omega) = \langle \tilde{\sigma}(\omega) \rangle$ is readily found to be

$$\langle \tilde{\sigma}(\omega) \rangle = \frac{1}{\omega - A + i\lambda B^2 + \tilde{K}(\omega)} \sigma(t_0) . \quad (7.27)$$

The effect of the finite correlation time, e.g., the deviation from the Wiener-Lévy limit, is accounted for by the operator

$$\bar{R}(\omega) = \gamma B \frac{1\sigma^2}{\omega - A + i\lambda B^2 + i1\gamma + \gamma B \frac{2\sigma^2}{\omega - A + i\lambda B^2 + 2i\gamma + \gamma B \frac{3\sigma^2}{\omega - A + i\lambda B^2 + 3i\gamma + \gamma B \dots}}} \gamma B \quad (7.28)$$

which indeed vanishes for $\gamma \rightarrow 0$, λ finite. In this limit, Eq. (7.27) is the Laplace transform of Eq. (7.8).

The explicit expression (7.27) provides the exact solution for situations where the initial state is non-stochastic and for cases where the solution is independent of the initial state. As an example from quantum optics, we mention that Eq. (7.27) with $\sigma(t_0) = 1$, $A = 0$ and $B = 1$ represents the laser spectral profile. Another example pertains to the long-time behavior of the solution $\langle \sigma(t) \rangle$. If there is any damping in the system, which might be caused by the stochastic fluctuations itself, then the solution for $t \gg t_0$ will become independent of the initial state. If we indicate by $\bar{\sigma}$ the solution $\langle \sigma(t) \rangle$ for $t \rightarrow \infty$, then $\bar{\sigma}$ obviously obeys the equation

$$(A - i\lambda B^2 - \bar{R}(0)) \bar{\sigma} = 0 \quad (7.29)$$

For the problem of atomic fluorescence in a strong laser field, this is the equation for the atomic steady-state density matrix, which determines the fluorescence yield. There, the solution $\bar{\sigma}$ of Eq. (7.29) is unique.

VIII. CONCLUSIONS

Solving a multiplicative stochastic process $\sigma(t)$ for its average is rarely feasible by analytical methods. This is mainly due to the finite

correlation time of the driving process $\phi(t)$, which prohibits the factorization of the average of a product into the product of the averages. Averages of a functional of $\phi(t)$ might factorize if the process is delta-correlated. For Markov processes, however, we can simulate a δ -correlation by the introduction of the marginal average $\zeta(\phi_0, t) = \langle \delta(\phi(t) - \phi_0) \alpha(t) \rangle$. The combination of the multiplication by $\delta(\phi(t) - \phi_0)$ and the Markov property of the probability distributions of $\phi(t)$ then gives rise to a factorization-like result for the formal expression for the average. Along the same lines as in a factorization assumption, we can now derive exact equations for $\zeta(\phi_0, t)$. In this paper we have studied Eq. (1.1), where we considered the stochastics of $\phi(t)$ to be given. The equation of motion for the marginal average turned out to be Eq. (6.4). The applicability of this equation was illustrated by some examples.

ACKNOWLEDGMENTS

This research was supported by the National Science Foundation under Grant CHE-8519053, the Air Force Office of Scientific Research (AFSC), United States Air Force, under Contract F49620-86-C-0009, and the Office of Naval Research. The United States Government is authorized to reproduce and distribute reprints for governmental purposes notwithstanding any copyright notation hereon.

REFERENCES

1. G. S. Agarwal, Phys. Rev. A 18, 1490 (1978).
2. A. T. George and S. N. Dixit, Phys. Rev. A 23, 2580 (1981).
3. H. F. Arnoldus and G. Nienhuis, J. Phys. B: At. Mol. Phys. (1986), in press.
4. R. F. Fox, J. Math. Phys. 13, 1196 (1972).
5. S. N. Dixit, P. Zoller and P. Lambropoulos, Phys. Rev. A 21, 1289 (1980).
6. P. Zoller, G. Alber and R. Salvador, Phys. Rev. A 24, 398 (1981).
7. R. I. Jackson and S. Swain, J. Phys. B: At. Mol. Phys. 15, 4375 (1982).
8. H. F. Arnoldus and G. Nienhuis, J. Phys. B: At. Mol. Phys. 16, 2325 (1983).
9. K. Wódkiewicz and J. H. Eberly, Phys. Rev. A 31, 2314 (1985).
10. A. Brissaud and U. Frisch, J. Quant. Spectr. Rad. Trans. 11, 1767 (1971).
11. A. Brissaud and U. Frisch, J. Math. Phys. 15, 524 (1974).
12. V. E. Shapiro and V. M. Loginov, Physica 91A, 563 (1978).
13. H. F. Arnoldus and G. Nienhuis, J. Phys. A: Math. Gen. 19, 1629 (1986).
14. B. W. Shore, J. Phys. Soc. Am. B 1, 183 (1984).
15. K. Wódkiewicz, B. W. Shore and J. H. Eberly, Phys. Rev. A 30, 2390 (1984).
16. P. Zoller and J. Cooper, Phys. Rev. A 28, 2310 (1983).
17. J. D. Cresser, W. H. Louisell, P. Meystre, W. Schleich and M. O. Scully, Phys. Rev. A 25, 2214 (1982).
18. N. G. van Kampen, Stochastic Processes in Physics and Chemistry (North-Holland, Amsterdam, 1981).

19. B. L. Stratonovich, Topics in the Theory of Random Noise (Gordon and Breach, New York, 1963).
20. R. F. Fox, Phys. Rev. A 33, 467 (1986).
21. N. G. van Kampen, Phys. Lett. 76A, 104 (1980).
22. A. Burshtein, Sov. Phys. JETP 21, 567 (1965).
23. J. J. Yeh and J. H. Eberly, Phys. Rev. A 24, 888 (1981).
24. H. Risken, The Fokker-Planck Equation: Methods of Solution and Applications (Springer, Berlin), in press.

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 413 800 N. Quincy Street Arlington, Virginia 22217	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 5042 Crane, Indiana 47522	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Washington, D.C. 20360	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
DTNSRDC Attn: Dr. G. Bosmajian Applied Chemistry Division Annapolis, Maryland 21401	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	Dr. David L. Nelson Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. G. A. Somorjai
Department of Chemistry
University of California
Berkeley, California 94720

Dr. J. Murday
Naval Research Laboratory
Surface Chemistry Division (6170)
455 Overlook Avenue, S.W.
Washington, D.C. 20375

Dr. J. B. Hudson
Materials Division
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Theodore E. Madey
Surface Chemistry Section
Department of Commerce
National Bureau of Standards
Washington, D.C. 20234

Dr. J. E. Demuth
IBM Corporation
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. M. G. Lagally
Department of Metallurgical
and Mining Engineering
University of Wisconsin
Madison, Wisconsin 53706

Dr. R. P. Van Duyne
Chemistry Department
Northwestern University
Evanston, Illinois 60637

Dr. J. M. White
Department of Chemistry
University of Texas
Austin, Texas 78712

Dr. D. E. Harrison
Department of Physics
Naval Postgraduate School
Monterey, California 93940

Dr. W. Kohn
Department of Physics
University of California, San Diego
La Jolla, California 92037

Dr. R. L. Park
Director, Center of Materials
Research
University of Maryland
College Park, Maryland 20742

Dr. W. T. Peria
Electrical Engineering Department
University of Minnesota
Minneapolis, Minnesota 55455

Dr. Keith H. Johnson
Department of Metallurgy and
Materials Science
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Dr. S. Sibener
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Arnold Green
Quantum Surface Dynamics Branch
Code 3817
Naval Weapons Center
China Lake, California 93555

Dr. A. Wold
Department of Chemistry
Brown University
Providence, Rhode Island 02912

Dr. S. L. Bernasek
Department of Chemistry
Princeton University
Princeton, New Jersey 08544

Dr. P. Lund
Department of Chemistry
Howard University
Washington, D.C. 20059

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. F. Carter
Code 6132
Naval Research Laboratory
Washington, D.C. 20375

Dr. Richard Colton
Code 6112
Naval Research Laboratory
Washington, D.C. 20375

Dr. Dan Pierce
National Bureau of Standards
Optical Physics Division
Washington, D.C. 20234

Dr. R. Stanley Williams
Department of Chemistry
University of California
Los Angeles, California 90024

Dr. R. P. Messmer
Materials Characterization Lab.
General Electric Company
Schenectady, New York 22217

Dr. Robert Gomer
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Ronald Lee
R301
Naval Surface Weapons Center
White Oak
Silver Spring, Maryland 20910

Dr. Paul Schoen
Code 5570
Naval Research Laboratory
Washington, D.C. 20375

Dr. John T. Yates
Department of Chemistry
University of Pittsburgh
Pittsburgh, Pennsylvania 15260

Dr. Richard Greene
Code 5230
Naval Research Laboratory
Washington, D.C. 20375

Dr. L. Kesmodel
Department of Physics
Indiana University
Bloomington, Indiana 47403

Dr. K. C. Janda
California Institute of Technology
Division of Chemistry and Chemical
Engineering
Pasadena, California 91125

Dr. E. A. Irene
Department of Chemistry
University of North Carolina
Chapel Hill, North Carolina 27514

Dr. Adam Heller
Bell Laboratories
Murray Hill, New Jersey 07974

Dr. Martin Fleischmann
Department of Chemistry
Southampton University
Southampton SO9 5NH
Hampshire, England

Dr. John W. Wilkins
Cornell University
Laboratory of Atomic and
Solid State Physics
Ithaca, New York 14853

Dr. Richard Smardzewski
Code 6130
Naval Research Laboratory
Washington, D.C. 20375

Dr. H. Tachikawa
Chemistry Department
Jackson State University
Jackson, Mississippi 39217

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. R. G. Wallis
Department of Physics
University of California
Irvine, California 92664

Dr. D. Ramaker
Chemistry Department
George Washington University
Washington, D.C. 20052

Dr. J. C. Hemminger
Chemistry Department
University of California
Irvine, California 92717

~~Dr. T. F. George
Chemistry Department
University of Rochester
Rochester, New York 14627~~

Dr. G. Rubloff
IBM
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. Horia Metiu
Chemistry Department
University of California
Santa Barbara, California 93106

Dr. W. Goddard
Division of Chemistry
California Institute of Technology
Pasadena, California 91125

Dr. J. T. Keiser
Department of Chemistry
University of Richmond
Richmond, Virginia 23173

Dr. R. W. Plummer
Department of Physics
University of Pennsylvania
Philadelphia, Pennsylvania 19104

Dr. E. Yeager
Department of Chemistry
Case Western Reserve University
Cleveland, Ohio 41106

Dr. N. Winograd
Department of Chemistry
Pennsylvania State University
University Park, Pennsylvania 16802

Dr. Roald Hoffmann
Department of Chemistry
Cornell University
Ithaca, New York 14853

Dr. A. Steckl
Department of Electrical and
Systems Engineering
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. G. H. Morrison
Department of Chemistry
Cornell University
Ithaca, New York 14853

Dr. P. Hansma
Physics Department
University of California
Santa Barbara, California 93106

Dr. J. Baldeschwieler
California Institute of Technology
Division of Chemistry
Pasadena, California 91125

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. J. E. Jensen
Hughes Research Laboratory
3011 Malibu Canyon Road
Malibu, California 90265

Dr. J. H. Weaver
Department of Chemical Engineering
and Materials Science
University of Minnesota
Minneapolis, Minnesota 55455

Dr. W. Goddard
Division of Chemistry
California Institute of Technology
Pasadena, California 91125

Dr. A. Reisman
Microelectronics Center of North Carolina
Research Triangle Park, North Carolina
27709

Dr. M. Grunze
Laboratory for Surface Science and
Technology
University of Maine
Orono, Maine 04469

Dr. J. Butler
Naval Research Laboratory
Code 6115
Washington D.C. 20375

Dr. L. Interante
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Irvin Heard
Chemistry and Physics Department
Lincoln University
Lincoln University, Pennsylvania 19352

Dr. K.J. Klaubunde
Department of Chemistry
Kansas State University
Manhattan, Kansas 66506

Dr. W. Knauer
Hughes Research Laboratory
3011 Malibu Canyon Road
Malibu, California 90265

Dr. C. B. Harris
Department of Chemistry
University of California
Berkeley, California 94720

Dr. F. Kutzler
Department of Chemistry
Box 5055
Tennessee Technological University
Cookeville, Tennessee 38501

Dr. D. DiLella
Chemistry Department
George Washington University
Washington D.C. 20052

Dr. R. Reeves
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

END

DTIC

9-86